

Heavy metals uptake pattern and accumulation in wheat

Nura ABDULLAHI

Department of Food Science and Technology, Aliko Dangote University of Science and Technology, Wudil, Kano State, Nigeria

Munir Abba DANDAGO

Department of Food Science and Technology, Aliko Dangote University of Science and Technology, Wudil, Kano State, Nigeria

Muhammad Sani GAMBO

Department of Crop Science, Aliko Dangote University of Science and Technology, Wudil, Kano State, Nigeria

Patrick Godwin IDAH

Department of Hospitality and Catering, Federal University of Health Sciences Otukpo, Benue State, Nigeria

Adamu Usman TSOHO

Department of Food Science and Technology, Aliko Dangote University of Science and Technology, Wudil, Kano State, Nigeria

Abdurrashid RILWAN

Department of Food Science and Technology, Aliko Dangote University of Science and Technology, Wudil, Kano State, Nigeria

The study aimed to determine the pattern of heavy metals (HMs) uptake and accumulation in wheat and compare the concentrations of the studied HMs in the wheat grain produced under the Kano River Irrigation Project with the permissible limits recommended by FAO/WHO. Wheat samples were collected from 10 different locations within the irrigation scheme between February and March 2023. The samples were prepared using standard laboratory procedures and HMs were determined using atomic absorption spectrophotometer. The accumulation of HMs in wheat grains is in the order of Cu>Zn>Ni>Cr>Co>Cd. Pb and Hg were not detected in wheat grains. The concentrations of Zn, Cd and Cr in the wheat grain samples exceeded the recommended permissible limits.

Keywords: Food contamination, Heavy metals toxicity, Nigeria, wheat contamination

INTRODUCTION

Wheat is a major staple crop, providing a significant portion of calories and protein to the world's population. It is the most widely cultivated crop, grown on millions of hectares annually (Erenstein et al., 2022). It is a highly nutritious grain that provides essential nutrients for human health. It is a rich source of carbohydrates, proteins, dietary fibre, lipids, vitamins, minerals, and phytochemicals (Wieser et al., 2020). Global wheat production exceeded 750 million metric tonnes (Mt)/annum, this accounts for about 20% of the calories in the human diet (Gooding, 2023). Nigeria's wheat consumption has been increasing over the years, driven by factors such as convenience, affordability, urbanization, and industrialization (Sadiq et al., 2020). Wheat is used for various purposes, including the production of bread, noodles, cookies, cakes, and breakfast cereals (Erenstein et al., 2022).

Heavy metals (HMs) infiltrate soil and water either through the natural process of rock weathering or as a result of human activities such as mining and the disposal of industrial waste (Bhagwat, 2019; Edelstein and Ben-Hur, 2018; Islam et al., 2019; Maqbool et al., 2019; Stefanović, et al., 2023a), abuse of agrochemicals and atmospheric deposit (Abdullahi et al., 2021). The presence of HMs in water and soil escalates the risk of food chain contamination (Taghipour and Jalali, 2019;

Saadati et al., 2020). The continuous release of hazardous HMs by various forms of human activities in quantities significantly surpassing those emitted by natural processes has created mounting concern regarding human health and environmental safety (Hu et al., 2018; Samiee et al., 2019; Vareda et al., 2019; Yuan et al., 2019). Excess HMs in the soil can directly alter its physical and chemical properties, decrease biological activity and diminish nutrient bioavailability (Kahkha et al., 2017). Production season is another important factor affecting soil HM behaviour and bioavailability (Salem et al., 2020).

Heavy metal contamination in wheat is of significant concern due to its potential impact on food safety, human health and crop quality. Their presence in food can cause complicated health issues that can lead to lead to death (Rai et al., 2019). Wheat plants are susceptible to accumulating high levels of toxic heavy metals such as cadmium lead and chromium (Zhou and Zheng, 2022). Various studies have highlighted the toxic effects of heavy metals on wheat, including impaired growth, yield losses, and biochemical responses (Zhou and Zheng, 2022). The presence of these metals in the soil can affect the density, composition, and physiological activities of microbiota, leading to a reduction in wheat production and posing risks to human and animal health (Rizvi et al., 2020; Zhou and Zheng, 2022). Additionally, the use of wastewater for irrigation has been associated with an increased risk of heavy metal accumulation in wheat grains (Mussarat et al., 2021). Various agrochemicals used for crop protection have also been reported to contaminate wheat with HMs and subsequently affect yield and nutritional qualities (Wołejko et al., 2017).

Various approaches have been studied to mitigate heavy metal uptake and phytotoxicity in wheat, including agronomic and biological methods (Dong et al., 2023), the use of plant growth-promoting rhizobacteria with heavy metal tolerance and plant growth-promoting potentials (Rizvi et al., 2020). The use of low-metal-accumulating or metal-toxicity-tolerant wheat varieties is considered the most efficient and eco-friendly approach to ensure food safety and protect human health (Saeed et al., 2023). However, the selection of the appropriate method depends on factors such as soil properties, type of metal, contamination level, and environmental conditions (Stefanović et al., 2023b). Understanding wheat's response to heavy metal stress and implementing management strategies to decrease metal uptake and accumulation is crucial for improving growth and grain quality (Rizvi et al., 2020).

The study aimed to determine the pattern of heavy metals (HMs) uptake and accumulation in wheat and compare the concentrations of the studied HMs in the wheat grain produced under the Kano River Irrigation Project with the permissible limits recommended by FAO/WHO.

METHODOLOGY

Sampling Location

The Kano River Irrigation Project (KRIP) covers approximately 62,000 hectares of land and only relies on the Tiga Dam and Ruwan Kanya Reservoir as its primary sources of irrigation water. This irrigation scheme is close to Kano City and extends across both sides of the Kano-Zaria expressway. Wheat samples were harvested at their optimal maturity and samples were collected between February and March 2023. Wheat grain and root samples were collected from 10 different locations across the irrigation scheme. The sampling locations are Gaiyare, Waire, Kadawa, Gafan, Makuntir, Kwalele Kura (K/Kudu), Kura (K/Gabas), Imawa I (Gada Mailotso) and Imawa II (Gonar Hajia).

Sample Preparation and HMs Determination

Harvested samples were separated into roots and grains. The grain samples were properly dried, trashed and adequately cleansed with deionised water. The root samples were soaked in deionised water before being extensively washed with an excess amount of water. The samples were then subjected to oven-drying until a constant weight was achieved. Then pulverized into a fine powder

and subsequently incinerated to ash using a muffle furnace at a temperature of 550 °C for 5 hours. The ash was mixed with concentrated HNO₃ and HClO₄ (3:1, v/v) in a heating digester. Each 1 g of the sample was mixed with 20 ml of the acid mixture. Then the acid digest was allowed to cool and filtered into 100 mL bottles, using Whatman filter paper and made up to mark with deionised water (Akinyele and Shokunbi, 2015). The concentrations of Zn, Pb, Cd, Ni, Co, Cu, Cr and Hg in the plant tissues were determined in triplicates using an atomic absorption spectrophotometer machine (Perkin Elmer PinAAcle 900H) and results were reported as Mean±SD.

Computation of Translocation Factor

A comparative analysis was conducted between the contents of heavy metals in the roots and those in the wheat grains to investigate the uptake patterns of Zn, Pb, Cd, Ni, Co, Cu, Cr, and Hg. The Translocation Factor (equation 1) was employed to examine the plant's capacity to transport the accumulated metals from the roots to the grains. TF was computed using the modified approach outlined by Zhuang et al. (2007).

$$TF = R_{conc.}/G_{conc.} \text{ Equation 1}$$

Where: R_{conc.} is the metal concentration in the root, and

G_{conc.} is the metal concentration in the grain.

RESULTS AND DISCUSSION

The results for HM contents of wheat grain are presented in Table 1. Mean concentrations for Zn, Cd, Ni, Co, Cu and Cr range between 1.640-6.713, 0.460-0.547, 2.012-4.053, ND-1.733, 3.873-5.353 and ND-11.127 mg/kg respectively. Pb and Hg were not detected in all the samples and Cr was only detected in samples collected from Gaiyare, Gafan, Makuntiri and Kwalele. The overall means for the sampling locations are 4.582, 0.509, 3.181, 0.623, 4.599 and 2.125 mg/kg for Zn, Cd, Ni, Co, Cu and Cr respectively. The overall mean concentration for Zn, Cd and Cr exceeded WHO-approved permissible limits.

The concentrations of Pb, Cd, Cr and Co reported by Madhi et al. (2020) are far beyond what was observed in this research. Also that of Zn in samples collected from Tianjin, China (Yu et al., 2016), Cr and Pb in pot experiment conducted in China (Fang et al., 2019), Zn and Pb in German wheat samples (Gramss, 2020), Zn and Pb in various samples collected from Pakistan (Al-Othman et al., 2016), Cr, Cu, Ni, Pb and Zn in samples produced at Shandong Province, China (Wan et al., 2022), Zn, Pb, Cr and Cu produced at Pancevo, Serbia (Stefanović et al., 2023b) Pb, Cu and Zn samples obtained from Xuzhou, China (Maqbool et al., 2019). Cu, Zn, Ni and Cd concentrations below what was observed in this study were also reported by Alemu et al. (2022) in wheat flour samples obtained from Amhara and Oromia, Ethiopia.

Concentrations below what was observed in this research were also reported by many researchers in wheat grains. These include the concentrations of Cd, Ni, Co, Cu and Cr in samples collected from various locations in Iran (Ghanati et al., 2019), Cd, Cu and Cr in samples collected from Tianjin, China (Yu et al., 2016), Cd, Zn and Cr in samples collected from Kosovo (Dreshaj et al., 2022), Ni, Zn and Cu in samples produced at Sistan, Iran (Kahkha et al., 2017), Zn in pot experiment conducted in China (Fang et al., 2019), Ni, Cd and Cr in German wheat samples (Gramss, 2020), Cd, Ni, Cu and Cr in various samples collected from Pakistan (Al-Othman et al., 2016), Cd in samples produced at Shandong Province, China (Wan et al., 2022), Cd and Cr in wheat produced at Xuzhou, China (Maqbool et al., 2019) and Cd produced at Pancevo, Serbia (Stefanović et al., 2023a). The variation in the concentrations of HM may be attributed to production location, cultural practices, wheat breed, soil properties, soil health, irrigation water quality and environmental safety.

The overall mean concentrations for Zn, Cd and Cr exceeded WHO-approved permissible limits. Madhi et al. (2020) reported Pb, Cd, Cr and Co concentrations above permissible limits in wheat samples collected from Basrah and Maysan. Hassan et al. (2013) also reported Cr and Ni concentrations above permissible limits in wheat samples collected from Abbottabad district, Pakistan. Wheat produced in Tianjin, China possessed Cd content above the permissible limit (Yu et al., 2016). On the other hand, Corguinha et al. (2015) reported As Cd and Pb concentrations below permissible limits in wheat grain produced in the states of Mato Grosso and Minas Gerais, Brazil.

Pb and Hg were not detected in all the samples. In the same way, Cd, Hg, and Pb were not detected in the wheat and barley grains imported to Egypt from some EU countries (Thabit et al., 2021). In contrast, Ghanati et al. (2019) detected both Pb and Hg in wheat and various wheat products. Similarly, Dreshaj et al. (2022) and Kahkha et al. (2017) detected Pb in wheat grains.

The results for the HM concentrations in the wheat root are presented in Table 2. Mean values for the concentrations of Zn, Cd, Ni, Co, Cu and Cr range between 2.787-6.460, 0.500-0.600, 2.720-3.987, 0.153-1.413, 3.747-5.260 and ND-5.647 mg/kg respectively. Similar to grain samples, Pb and Hg were also not detected in all the root samples. Cr was only detected in samples collected from Gaiyare, Kadawa, Makuntiri, Kwalele and Imawa I (Gada Mailotso). Contrary to what was observed in the grains, Cr was detected in the root samples collected from Kadawa and Imawa I. The overall means for the concentrations of Zn, Cd, Ni, Co, Cu and Cr are 4.985, 0.528, 3.359, 0.675, 4.414 and 2.288 mg/kg respectively. Higher mean values for Zn, Cd, Ni and Co were found in root samples while higher mean values for Cu and Cr were found in the grain samples. Yu et al. (2016) reported Cu, Pb, Zn and Ni concentrations above what was observed in this study in the root of wheat produced at Tianjin. Also, the concentration of Pb in the root of various samples collected from Pakistan (Al-Othman et al., 2016). Concentrations Cd, Ni, Cu, Zn and Cr below what was observed in this research were reported by Al-Othman et al. (2016) in the root of wheat produced in Pakistan.

Translocation factor

Calculated values for translocation factors of wheat are presented in Table 3. A Root:Grain metal concentration value >1 signifies higher accumulation in the roots while values <1 indicate accumulation in the grains. The overall mean values for TF show higher accumulations of Zn, Cd, Ni and Co in the root and higher accumulation of Cu and Cr in the grain. Similar to this finding, Al-Othman et al. (2016) also reported higher concentrations of HMs in the wheat root than in the grain.

Accumulation in the Root: $R_{conc.}/G_{conc.} > 1$

Accumulation in the Grains: $R_{conc.}/G_{conc.} < 1$

CONCLUSION

The accumulation of HMs in wheat grains is in the order of $Cu > Zn > Ni > Cr > Co > Cd$. Pb and Hg were not detected in wheat grains. The concentrations of Zn, Cd and Cr in the wheat grain samples exceeded the recommended permissible limits. Controlling HMs contamination in wheat requires production on good quality soil and the use of only good quality water for irrigation. The use of low-metal-accumulating or metal-toxicity-tolerant wheat varieties is considered the most efficient and eco-friendly approach. Microbiological remediation strategies, such as the use of metal-tolerant plant growth-promoting rhizobacteria, have shown promise in enhancing wheat production in metal-stressed soil.

Funding

The research was fully funded by the Hedija Jama'are River Basin Development Authority.

REFERENCES

- Abdullahi N., Igwe E.C., Dandago M.A. (2021). Heavy metals contamination through consumption of contaminated food crops. *Moroccan Journal of Agricultural Sciences*, 2: 52-60.
- Akinyele I.O., Shokunbi O.S. (2015). Comparative analysis of dry ashing and wet digestion methods for the determination of trace and heavy metals in food samples. *Food Chemistry*, 173: 682-684.
- Al-Othman Z.A., Ali R., Al-Othman A.M., Ali J., A. Habila M. (2016). Assessment of toxic metals in wheat crops grown on selected soils, irrigated by different water sources. *Arabian Journal of Chemistry*, 9: 1555-1562.
- Alemu W.D., Bulta A.L., Doda M.B., Kanido C.K. (2022). Levels of selected essential and non-essential metals in wheat (*Triticum aestivum*) flour in Ethiopia. *Journal of Nutritional Science*, 11: 1-9.
- Bhagwat V.R. (2019). Safety of water used in food production. In R. L. Singh and S. Mondal (Eds.), *Food Safety and Human Health*: 219-247. Elsevier Inc.
- Chiroma T.M., Ebewele R.O., Hymore F. (2014). Comparative assessment of heavy metal levels in soil, vegetables and urban grey waste water used for irrigation in Yola And Kano. *International Refereed Journal of Engineering and Science*, 3: 1-09.
- Corguinha A.P.B., Souza G.A. de, Gonçalves V.C., Carvalho C. de A., Lima W.E.A. de, Martins F.A.D., Yamanaka C.H., Francisco E.A.B., Guilherme L.R.G. (2015). Assessing arsenic, cadmium, and lead contents in major crops in Brazil for food safety purposes. *Journal of Food Composition and Analysis*, 37: 143-150.
- Dong W., Hu T., Zhang Q., Deng F., Wang M., Kong J., Dai Y. (2023). Prediction of food safety risk level of wheat in china based on pyraformer neural network model for heavy metal contamination. *Foods*, 12(9): 1-19.
- Dreshaj A., Millaku B., Gashi A., Elezaj E., Kuqi, B. (2022). Concentration of toxic metals in agricultural land and wheat culture (*Triticum aestivum* L.). *Journal of Ecological Engineering*, 23: 18-24.
- Edelstein M., Ben-Hur M. (2018). Heavy metals and metalloids: Sources, risks and strategies to reduce their accumulation in horticultural crops. *Scientia Horticulturae*, 234: 431-444.
- Erenstein O., Jaleta M., Abdul Mottaleb K., Sonder K., Donovan J., Braun H.J. (2022). Global trends in wheat production, consumption and trade. *Wheat Improvement: Food Security in a Changing Climate*, 47-66.
- Fang A., Dong J., Zhang R. (2019). Simulation of heavy metals migration in soil-wheat system of mining area. *International Journal of Environmental Research and Public Health*, 16: 1-14.
- Ghanati K., Zayeri F., Hosseini H. (2019). Potential health risk assessment of different heavy metals in wheat products. *Iranian Journal of Pharmaceutical Research*, 18: 2093-2100.
- Gooding M. J. (2023). Wheat. In *ICC Handbook of 21st century cereal science and technology* (pp. 121-130). Academic Press.

Gramss G. (2020). Control of heavy metals from barley and wheat grains during malting and brewing. *Advances in Nutrition and Food Science*, 2020: 1-9.

Hassan N.U., Mahmood Q., Waseem A., Irshad M., Faridullah, Pervez A. (2013). Assessment of heavy metals in wheat plants irrigated with contaminated wastewater. *Polish Journal of Environmental Studies*, 22: 115-123.

Hu W., Wang H., Dong L., Huang B., Borggaard O.K., Bruun Hansen H.C., He Y., Holm P.E. (2018). Source identification of heavy metals in peri-urban agricultural soils of southeast China: An integrated approach. *Environmental Pollution*, 237: 650-661.

Islam M.S., Ahmed M.K., Al-mamun M.H., Islam S.M. A. (2019). Sources and ecological risks of heavy metals in soils under different land uses in Bangladesh. *Pedosphere*, 29: 665-675.

Kahkha M.R.R., Bagheri S., Noori R., Piri J., Javan S. (2017). Examining total concentration and sequential extraction of heavy metals in agricultural soil and wheat. *Polish Journal of Environmental Studies*, 26: 2021-2028.

Madhi Q.H., Abass M.H., Matrood, A.A.A. (2020). Heavy metals pollution of wheat fields (soil and leaves) sampled from Basrah and Maysan Provinces. *Periodico Tche Quimica*, 17: 315-326.

Maqbool A., Xiao X., Wang H., Bian Z., Akram M. W. (2019). Bioassessment of heavy metals in wheat crop from Soil and dust in a coal mining area. *Pollution*, 5: 323-337.

Mussarat M., Jamal W.A., Muhammad D., Ahmad M., Saleem A., Khan S., Aman F., Bibi H., Shah W.A., Dawar K., Ul Akbar N., Mian I.A., Waheed M., Ali I., Zia A., Malik, W. (2021). Risk of heavy metals accumulation in soil and wheat grains with waste water irrigation under different NPK levels in alkaline calcareous soil. *PLoS ONE*, 16: 1-15.

Ogundele D., Adio A., Oludele O. (2015). Heavy metal concentrations in plants and soil along heavy traffic roads in north central Nigeria. *Journal of Environmental and Analytical Toxicology*, 5: 6-10.

Rai P.K., Lee S.S., Zhang M., Tsang Y.F., Kim K.H. (2019). Heavy metals in food crops: Health risks, fate, mechanisms, and management. *Environment International*, 125: 365-385.

Rizvi A., Zaidi A., Ameen F., Ahmed B., Alkahtani M.D. F., Khan M.S. (2020). Heavy metal induced stress on wheat: Phytotoxicity and microbiological management. *RSC Advances*, 10: 38379-38403.

Saadati M., Soleimani M., Sadeghsaba M., Hemami M. R. (2020). Bioaccumulation of heavy metals (Hg, Cd and Ni) by sentinel crab (*Macrophthalmus depressus*) from sediments of Mousa Bay, Persian Gulf. *Ecotoxicology and Environmental Safety*, 191: 1-7.

Sadiq S., Sngh I.P., Ahmad M.M. (2020). Domestic consumption pattern of cereal commodities in Nigeria. *Alnteri Zirai Bilimler Dergisi*, 35: 120-126.

Saeed M., Quraishi U.M., Malik, R.N. (2023). Advancement in mitigating the effects of heavy metal toxicity in wheat. *Abiotic Stresses in Wheat: Unfolding the Challenges*, 313-327.

Salem M.A., Bedade D.K., Al-Ethawi L., Al-waleed S. M. (2020). Assessment of physiochemical properties and concentration of heavy metals in agricultural soils fertilized with chemical fertilizers. *Heliyon*, 6: 1-20.

Samiee F., Vahidinia A., Taravati Javad M., Leili M. (2019). Exposure to heavy metals released to the environment through breastfeeding: A probabilistic risk estimation. *Science of the Total Environment*, 650: 3075-3083.



Stefanović V.M., Nikolić S.R., Matković Stojšin M., Majstorović H., Petreš M., Cvikić D., Racić G. (2023a). Soil-to-wheat transfer of heavy metals depending on the distance from the industrial zone. *Agronomy*, 13:1–23.

Stefanović V.M., Stanisavljević D., Bačić J., Brković P., Pavlović M., Matković- Stojšin M. (2023b). Influence of Heavy Metals on Wheat Stem Development. 1st International Symposium on Biotechnology, 55–62.

Taghipour M., Jalali M. (2019). Impact of some industrial solid wastes on the growth and heavy metal uptake of cucumber (*Cucumis sativus* L.) under salinity stress. *Ecotoxicology and Environmental Safety*, 182: 1–11.

Thabit T.M.A.M., Shokr S.A., Elgeddawy D.I.H., El-Naggar M.A.H. (2021). Determination of heavy metals in wheat and barley grains using ICP-MS/MS. *Journal of AOAC International*, 103: 1277–1281.

Vareda J.P., Valente A.J.M., Durães L. (2019). Assessment of heavy metal pollution from anthropogenic activities and remediation strategies: A review. *Journal of Environmental Management*, 246: 101–118.

Wan F., Jiang N., Yu L., Zang K., Liu S., He W., Hu Z., Fan H., Li H., Wang H., Pan H., Yang Q., Lou Y., Zhuge Y. (2022). Heavy metal ecological-health risk assessment under wheat-maize rotation system in a high geological background area in eastern China. *Scientific Reports*, 12: 1–12.

Wieser H., Koehler P., Scherf K.A. (2020). Nutritional value of wheat. In *Wheat - An Exceptional Crop* (pp. 133–148). Woodhead Publishing.

Wolejko E., Łozowicka B., Kaczyński P., Konecki R., Grobela M. (2017). The influence of chemical protection on the content of heavy metals in wheat (*Triticum aestivum* L.) growing on the soil enriched with granular sludge. *Environmental Monitoring and Assessment*, 189: 1–12.

Yu X., Wang Z., Lynn A., Cai J., Huangfu Y., Geng Y., Tang J., Zeng X. (2016). Heavy metals in wheat grown in sewage irrigation: A distribution and prediction model. *Polish Journal of Environmental Studies*, 25: 413–418.

Yuan Z., Luo T., Liu X., Hua H., Zhuang Y., Zhang X., Zhang L., Zhang Y., Xu W., Ren J. (2019). Tracing anthropogenic cadmium emissions: From sources to pollution. *Science of the Total Environment*, 676: 87–96.

Zhou M., Zheng S. (2022). Multi-omics uncover the mechanism of wheat under heavy metal stress. *International Journal of Molecular Sciences*, 23: 1–16.

Zhuang P., Yang Q.W., Wang H.B., Shu W.S. (2007). Phytoextraction of heavy metals by eight plant species in the field. *Water, Air, and Soil Pollution*, 184: 235–242.

References